

5. Übungsblatt
Theoretische Physik 6: WS 2014/15
Dozent: Prof. M. Vanderhaeghen

24.11.2014

Exercise 1 (40 points): Real Klein-Gordon field

Using the normal mode expansion of the real Klein-Gordon field,

$$\phi(\vec{x}, t) = \sum_{\vec{k}} \left(\frac{\hbar c^2}{2\omega_k L^3} \right)^{1/2} \left[a(\vec{k}) e^{-ik \cdot x} + a^\dagger(\vec{k}) e^{ik \cdot x} \right],$$

and the equal-time commutation relations:

$$\begin{aligned} [\phi(\vec{x}, t), \phi(\vec{x}', t)] &= 0, \\ [\dot{\phi}(\vec{x}, t), \dot{\phi}(\vec{x}', t)] &= 0, \\ [\phi(\vec{x}, t), \dot{\phi}(\vec{x}', t)] &= i\hbar c^2 \delta^{(3)}(\vec{x} - \vec{x}'). \end{aligned}$$

Show that:

(a) (20 points) the creation and annihilation operators satisfy the following commutation relations:

$$\begin{aligned} [a(\vec{k}), a(\vec{k}')] &= 0, \\ [a^\dagger(\vec{k}), a^\dagger(\vec{k}')] &= 0, \\ [a(\vec{k}), a^\dagger(\vec{k}')] &= \delta_{\vec{k}, \vec{k}'}; \end{aligned}$$

(b) (10 points) the Hamiltonian $H = \int d^3x \frac{1}{2} \left[\frac{1}{c^2} \dot{\phi}^2 + (\vec{\nabla} \phi)^2 + \mu^2 \phi^2 \right]$ takes the form:

$$H = \sum_{\vec{k}} \hbar \omega_k \left(a^\dagger(\vec{k}) a(\vec{k}) + \frac{1}{2} \right),$$

(c) (10 points) the momentum $\vec{P} = - \int d^3x \frac{1}{c^2} \dot{\phi} \vec{\nabla} \phi$ takes the form:

$$\vec{P} = \sum_{\vec{k}} \hbar \vec{k} \left(a^\dagger(\vec{k}) a(\vec{k}) + \frac{1}{2} \right),$$

Exercise 2 (60 points): Complex Klein-Gordon field

The complex Klein-Gordon field is used to describe charged bosons. Its Lagrangian is given by

$$\mathcal{L} = (\partial_\mu \phi^\dagger)(\partial^\mu \phi) - \mu^2 \phi^\dagger \phi, \quad (1)$$

where the field ϕ has the following normal mode expansion

$$\phi(\vec{x}, t) = \sum_{\vec{k}} \left(\frac{\hbar c^2}{2\omega_k L^3} \right)^{1/2} \left[a(\vec{k}) e^{-ik \cdot x} + b^\dagger(\vec{k}) e^{ik \cdot x} \right]$$

and satisfies the equal-time commutation relations:

$$\begin{aligned} [\phi(\vec{x}, t), \Pi_\phi(\vec{x}', t)] &= i\hbar \delta^{(3)}(\vec{x} - \vec{x}'), \\ [\phi^\dagger(\vec{x}, t), \Pi_{\phi^\dagger}(\vec{x}', t)] &= i\hbar \delta^{(3)}(\vec{x} - \vec{x}'); \end{aligned}$$

In the following, you can conveniently consider the fields ϕ and ϕ^\dagger as independent.

(a) (15 points) Show that (1) is equivalent to the Lagrangian of two independent real scalar fields with same mass and satisfying the standard equal-time commutation relations. *Hint:* Decompose the complex field in real components $\phi = \frac{1}{\sqrt{2}} (\phi_1 + i\phi_2)$.

(b) (15 points) Write down the conjugate momentum fields Π_ϕ and Π_{ϕ^\dagger} in terms of ϕ and ϕ^\dagger , and derive the equal-time commutation relations of a , a^\dagger , b and b^\dagger .

(c) (15 points) Show that (1) is invariant under any global phase transformation of the field $\phi \rightarrow \phi' = e^{-i\alpha} \phi$ with α real. Write down the associated conserved Noether current J^μ and express the conserved charge $Q = \int d^3x J^0$ in terms of creation and annihilation operators.

(d) (15 points) Compute the commutators $[Q, \phi]$ and $[Q, \phi^\dagger]$. Using these commutators and the eigenstates $|q\rangle$ of the charge operator Q , show that the field operators ϕ and ϕ^\dagger modify the charge of the system. How would you interpret the operators a , a^\dagger , b and b^\dagger ?