

Supporting Information

Micro-site conditions affect Fennoscandian forest growth

Claudia Hartl¹, Elisabeth Dũthorn¹, Ernesto Tejedor², Andreas J. Kirchhefer³, Mauri Timonen⁴, Steffen Holzkamper⁵, Ulf Bũntgen⁶⁻⁸, and Jan Esper¹

¹ Department of Geography, Johannes Gutenberg University, Mainz, German; ² Department of Atmospheric and Environmental Sciences, University at Albany (SUNY), Albany (New York), USA; ³ Dendrookologen A.J. Kirchhefer, Tromso, Norway; ⁴ Natural Resources Institute Finland, Etelaranta 55, Rovaniemi, Finland; ⁵ Department of Physical Geography, Stockholm University, Stockholm, Sweden; ⁶ Department of Geography, University of Cambridge, UK; ⁷ Swiss Federal Research Institute WSL, Birmensdorf, Switzerland; ⁸ Global Change Research Centre AS CR, Brno, Czech Republic

Corresponding author: Claudia Hartl (c.hartl@geo.uni-mainz.de)

-
- Fig. S1.** Cambial age aligned mean micro-site growth rates. Red = dry and blue = moist micro-sites.
- Fig. S2.** Monthly climate response of the micro-site chronologies with E-OBS v19.0 climate data (Cornes et al., 2018) and the 1950-2006 period.
- Fig. S3.** July temperature response over time (25-years running correlations) at each micro site for the 1950 to 2006 period.
-




Fig. S1. Cambial age aligned mean micro-site growth rates. Red = dry and blue = moist micro-sites.




Fig. S2. Monthly climate response of the micro-site chronologies with E-OBS v19.0 climate data (Cornes et al., 2018) and the 1950-2006 period.




Fig. S3. July temperature response over time (25-years running correlations) at each micro site for the 1950 to 2006 period.